

Crystal Data: Monoclinic. *Point Group:* 2/m. As aggregates (to 1.5 cm) of acicular crystals to 1.0 mm, and as separate elongated columnar, flattened-prismatic crystals to 1 cm with rectangular or rhombic sections.

Physical Properties: *Cleavage:* Very good on {100}. *Tenacity:* Brittle. *Fracture:* Uneven, conchoidal. VHN = 547–659, 569 average (100 g load). Hardness = ~5 D(meas.) = 3.44(2) D(calc.) = 3.475

Optical Properties: Transparent. *Color:* Colorless, white. *Streak:* White. *Luster:* Vitreous. *Optical Class:* Biaxial (+). $\alpha = 1.662(2)$ $\beta = 1.666(2)$ $\gamma = 1.685(5)$ $2V(\text{meas.}) = 50(3)^\circ$ $2V(\text{calc.}) = 49.7^\circ$ *Dispersion:* Medium, $v > r$. Nonpleochroic.

Cell Data: *Space Group:* $P2_1/c$. $a = 7.3934(5)$ $b = 5.6347(4)$ $c = 18.713(1)$ $\beta = 101.415(2)^\circ$ $Z = 2$

X-Ray Diffraction Pattern: Darai-Pioz alkaline massif, Tien Shan mountains, Tajikistan. 3.057 (100), 2.688 (28), 9.18 (24), 2.929 (17), 3.559 (15), 2.293 (15), 2.783 (14)

Chemistry:

	(1)		(1)
WO_3	0.41	Pr_2O_3	0.61
Ta_2O_5	0.15	Ce_2O_3	3.18
Nb_2O_5	2.74	La_2O_3	0.56
UO_2	0.22	Y_2O_3	6.82
TiO_2	8.32	SrO	0.35
SiO_2	29.51	MnO	0.28
Dy_2O_3	1.35	CaO	25.53
Gd_2O_3	1.58	Na_2O	7.98
Sm_2O_3	0.99	F	6.02
Nd_2O_3	3.34	<u>-O=F</u>	<u>2.53</u>
		Total	97.41

(1) Darai-Pioz alkaline massif, Tien Shan mountains, Tajikistan; average electron microprobe analysis supplemented by IR spectroscopy; corresponds to $\text{Na}_{2.11}(\text{Ca}_{3.74}\text{Sr}_{0.03}\text{Mn}_{0.03})_{\Sigma=3.80}(\text{Y}_{0.50}\text{Nd}_{0.16}\text{Ce}_{0.16}\text{Gd}_{0.07}\text{Dy}_{0.06}\text{Sm}_{0.05}\text{Pr}_{0.03}\text{La}_{0.03}\text{U}^{4+})_{\Sigma=1.07}(\text{Ti}_{0.85}\text{Nb}_{0.17}\text{W}^{6+}_{0.01}\text{Ta}_{0.01})_{\Sigma=1.04}(\text{Si}_{4.03}\text{O}_{14})\text{O}_{1.40}\text{F}_{2.60}$.

Mineral Group: Seidozerite supergroup, rinkite group.

Occurrence: In a pegmatite fragment from an alkaline massif collected from the moraine of the Darai-Pioz glacier.

Association: Quartz, aegirine, microcline, neptunite, pectolite, calcite, eudialyte-group minerals, fluorite, titanite, turkestanite, kupletskite, galena, albite, pyrochlore-group minerals.

Distribution: From the Darai-Pioz alkaline massif, Tien Shan mountains, Tajikistan.

Name: Identifies the Y-analog of *rinkite-(Ce)*.

Type Material: A.E. Fersman Mineralogical Museum, RAS, Moscow, Russia (5043/1).

References: (1) Pautov, L.A., A.A. Agakhanov, V.Y. Karpenko, Y.A. Uvarova, E. Sokolova, and F.C. Hawthorne (2019) Rinkite-(Y), $\text{Na}_2\text{Ca}_4\text{YTi}(\text{Si}_2\text{O}_7)_2\text{OF}_3$, a seidozerite-supergroup TS-block mineral from the Darai-Pioz alkaline massif, Tien-Shan mountains, Tajikistan: Description and crystal structure. *Mineral. Mag.*, 83, 373–380.