

**Crystal Data:** Hexagonal. *Point Group:* 6/*m*. Crystals acicular, prismatic, to 3 mm, in tufted groups, spherical rosettes.

**Physical Properties:** Hardness = n.d. D(meas.) = 3.65(5) D(calc.) = 3.62(1)

**Optical Properties:** Semitransparent. *Color:* Grass-green to dull green, yellowish green to intense bluish green, rarely nearly colorless.

*Optical Class:* Uniaxial (+).  $\omega = 1.715(2)$   $\varepsilon = 1.795(2)$

**Cell Data:** *Space Group:* P6<sub>3</sub>/*m*.  $a = 13.586(4)$   $c = 5.931(5)$   $Z = 2$

**X-ray Powder Pattern:** n.d. ; presumably very similar to agardite-(Y).

**Chemistry:** (1) Red Cloud fluorite mine, New Mexico, USA; analysis not given, by electron microprobe, average of analyses on four crystals, (OH)<sup>1-</sup> calculated for charge balance; stated to correspond to (Al<sub>0.36</sub>La<sub>0.29</sub>Ce<sub>0.21</sub>Nd<sub>0.08</sub>Pr<sub>0.03</sub>Y<sub>0.02</sub>Gd<sub>0.02</sub>Sm<sub>0.01</sub>)<sub>Σ=1.02</sub>(Cu<sub>5.50</sub>Ca<sub>0.42</sub>Pb<sub>0.06</sub>Fe<sub>0.02</sub>Zn<sub>0.01</sub>)<sub>Σ=6.01</sub> [(AsO<sub>4</sub>)<sub>2.74</sub>(SiO<sub>4</sub>)<sub>0.25</sub>(VO<sub>4</sub>)<sub>0.02</sub>(SO<sub>4</sub>)<sub>0.02</sub>]<sub>Σ=3.03</sub>(OH)<sub>5.76</sub>·3H<sub>2</sub>O.

**Mineral Group:** Mixite group; Nd and Ce have also been noted as dominant rare-earth elements, but the corresponding species have not been fully described.

**Occurrence:** In small amounts in the oxidized zone of hydrothermal mineralized breccia and polymetallic mineral deposits (Red Cloud mines, New Mexico, USA).

**Association:** Fluorite, bastnaesite, barite, quartz (Red Cloud fluorite mine, New Mexico, USA); chrysocolla, malachite, azurite, mimetite, vanadinite, conichalcite, wulfenite, mottramite, cerussite, quartz (Red Cloud copper mine, New Mexico, USA); smithsonite, aurichalcite, hydrozincite, azurite, cuprian adamite, calcite, chrysocolla, zincaluminite, gibbsite (Kamariza mine, Greece).

**Distribution:** From the Red Cloud fluorite and copper mines, Gallinas district, Lincoln Co., New Mexico, USA. In the Kamariza mine, Laurium, Greece (Ce > La in part). In England, from Wheal Alfred, Phillack, Cornwall (Nd-rich); at the Brandy Gill and Deer Hills mines, Caldbeck Fells, Cumbria. At the Clara Mine, near Oberwolfach, and other localities in the Black Forest, Germany (Ce > La in part). From the Sa Duchessa mine, near Iglesias, Sardinia, Italy (Nd-rich in part, some with Ce > La).

**Name:** By analogy to agardite-(Y), with its rare-earth content dominated by lanthanum.

**Type Material:** National Museum of Natural History, Washington, D.C., USA, 148987, 148988.

**References:** (1) Modreski, P.J. (1983) Agardite-(La), a chemically complex rare-earth arsenate from the Gallinas district, Lincoln Co., New Mexico. In: Anthony, J.W., Ed., Oxidation mineralogy of base metal deposits: Fifth Joint Mineralogical Society of America - Friends of Mineralogy Symposium, Tucson, Arizona. (2) Fehr, T. and R. Hochleitner (1984) Agardite-La, ein neues Mineral von Lavrion, Griechenland. *Lapis*, 9(1), 22, 37 (in German). (3) (1985) *Amer. Mineral.*, 70, 871 (abs. ref. 2). (4) Golubev A.M., E. Brücher, A. Schulz, R.K. Kremer, and R. Glaum (2020) La- and Lu-agardite - preparation, crystal structure, vibrational and magnetic properties. *Zeitschrift für Naturforschung B Chemical Science*, 75, 191-199.