

Crystal Data: Monoclinic, pseudo-orthorhombic. *Point Group:* $2/m$. As water-worn, bean-shaped pebbles, to 10 cm, with polycrystalline or radial-fibrous structure; also as curved and striated crystals, diamond-shaped or rectangular, lining pockets in the pebbles. *Twinning:* As pseudohexagonal multiple contact twins, chrysoberyl-like.

Physical Properties: *Cleavage:* {100}, perfect. *Fracture:* Uneven. Hardness = 9 VHN = 1605 (100 g load). $D(\text{meas.}) = 4.78\text{--}5.29$; 5.46 for a crystal. $D(\text{calc.}) = 5.26$

Optical Properties: Transparent to translucent. *Color:* Tan, cream, orange-brown, brown; crystals colorless, tan, pale violet; yellow internal reflections in transmitted light. *Luster:* Adamantine.

Optical Class: Biaxial (-). *Pleochroism:* Weak. *Dispersion:* $r > v$. $\alpha = 1.81(1)$ $\beta = 1.87(1)$ $\gamma = 1.92(1)$ $2V(\text{meas.}) = \text{Large}$.

Cell Data: *Space Group:* $C2/m$. $a = 9.406(6)$ $b = 11.541(8)$ $c = 4.410(3)$ $\beta = 90.94(3)^\circ$ $Z = 2$

X-ray Powder Pattern: Paramirim region, Brazil.

3.241 (100), 3.194 (100), 2.156 (80), 4.712 (70), 2.457 (70), 2.411 (70), 1.648 (70)

Chemistry:	(1)	(2)	(3)
WO_3	1.20	0.3	
Sb_2O_5	57.28	61.9	64.00
SiO_2	1.03	3.5	
TiO_2		0.2	
Al_2O_3	35.37	30.6	33.62
Fe_2O_3	1.04	1.3	
BeO	0.75	0.75	
MgO		0.1	
H_2O	2.77	0.85	2.38
Total	99.44	99.50	100.00

(1) Paramirim region, Brazil; by electron microprobe, average of four analyses, total Fe as Fe_2O_3 , H_2O by the Penfield method; corresponds to $\text{Al}_{5.49}\text{Be}_{0.24}\text{Si}_{0.14}\text{Fe}_{0.10}^{3+}\text{W}_{0.04}\text{Sb}_{2.80}\text{O}_{14}(\text{OH})_{2.44}$.

(2) Do. (3) $\text{Al}_5\text{Sb}_3\text{O}_{14}(\text{OH})_2$.

Occurrence: In potholes as stream-worn pebbles and as residual concentrates above weathered volcanic rock.

Association: Quartz, andalusite, kyanite, diaspore, phenakite, eskolaite, cassiterite, gold.

Distribution: About eight km southeast of the village of Paramirim das Crioulas, and elsewhere in the region around Rio do Picos das Almas, Serra das Almas, south-central Bahia, Brazil.

Name: For the occurrence in Bahia, Brazil.

Type Material: Harvard University, Cambridge, Massachusetts, 119088; National Museum of Natural History, Washington, D.C., USA, 133875, 135922.

References: (1) Moore, P.B., C. do P. Barbosa, and R.V. Gaines (1978) Bahianite, $\text{Sb}_3\text{Al}_5\text{O}_{14}(\text{OH})_2$, a new species. Mineral. Mag., 42, 179–182. (2) (1979) Amer. Mineral., 64, 464 (abs. ref. 1). (3) Moore, P.B. and T. Araki (1976) Bahianite, $\text{Al}_6\text{Sb}_3^{5+}\text{O}_{14}(\text{O}, \text{OH})_2$, a novel hexagonal close-packed oxide structure. Neues Jahrb. Mineral., Abh., 126, 113–125. (4) Cassedanne, J.P. (1985) Bahianite from Brazil. Mineral. Record, 16, 111–115.

All rights reserved. No part of this publication may be reproduced, stored in a retrieval system or transmitted in any form or by any means, electronic, mechanical, photocopying, recording, or otherwise without the prior written permission of Mineral Data Publishing.