


Birnessite

©2001-2005 Mineral Data Publishing, version 1

Crystal Data: Monoclinic, pseudo-hexagonal. *Point Group:* $2/m$. Rarely in platelets, to 50 μm ; commonly extremely finely crystalline, spherulitic, cellular.**Physical Properties:** Hardness = 1.5 D(meas.) = 3.0 D(calc.) = [3.4]**Optical Properties:** Nearly opaque. *Color:* Black; dark brown in transmitted light.*Luster:* Dull.*Optical Class:* Pseuduniaxial (-); identification by optical properties is impossible. $\omega = \sim 1.73$
 $\epsilon = \sim 1.69$ **Cell Data:** *Space Group:* $C2/m$ (synthetic $\approx \text{Na}_{0.58}(\text{Mn}_{1.42}^{4+}\text{Mn}_{0.58}^{3+})_{\Sigma=2.00}\text{O}_4 \cdot 1.5\text{H}_2\text{O}$).
 $a = 5.175(1)$ $b = 2.850(1)$ $c = 7.337(3)$ $\beta = 103.18(2)^\circ$ $Z = 1$ **X-ray Powder Pattern:** Marine nodule, Caribbean; all reflections are typically very broad.
7.08 (100), 2.333 (43), 1.711 (29), 3.547 (28), 2.031 (24), 2.468 (17), 1.426 (17)

Chemistry:	(1)	(2)	(1)	(2)
SiO ₂	0.9		CaO	0.39
MnO ₂	75.8	61.40	Na ₂ O	1.9
Mn ₂ O ₃		18.58	K ₂ O	1.8
FeO	0.55		Cl	0.1
CoO	0.14		H ₂ O	[10.89]
NiO	0.80		S	0.2
CuO	0.33		rem.	
MgO	6.2		Total	[100.0]
				100.00

(1) Marine nodule, Caribbean; by electron microprobe, total Mn as MnO₂, H₂O by difference.(2) $\text{Na}_{0.50}(\text{Mn}_{1.50}^{4+}\text{Mn}_{0.50}^{3+})_{\Sigma=2.00}\text{O}_4 \cdot 1.5\text{H}_2\text{O}$.**Occurrence:** A major manganese-bearing mineral of many soils; a common alteration product of manganese-rich mineral deposits; a component of bacterially-precipitated manganese oxides; an important constituent of "desert varnish" and marine manganese nodules.**Association:** Rhodonite, rhodochrosite, tephroite, spessartine, alleghanyite, cummingtonite (Cummington, Massachusetts, USA); manganese and iron oxides, calcium carbonate (marine nodules).**Distribution:** Increasingly recognized as a common manganese oxide mineral. From Birness, Aberdeenshire, Scotland. In the Treburland mine, Altarnun, and the Penberthy Croft mine, St. Hilary, Cornwall, England. On Groote Eylandt, Northern Territory, and near Noarlunga, Mount Lofty Ranges, South Australia. At the Ioi mine, Ritta, Shiga Prefecture, Japan. In the USA, at Sterling Hill, Sussex Co., New Jersey; from Cummington, Hampshire Co., Massachusetts; in the Kramer borate deposit, Boron, Kern Co., California; at Silver Cliff, Custer Co., Colorado; and elsewhere. From Mont St. Hilaire, Quebec, Canada. A number of oceanic sea floor and sea mount occurrences are now known.**Name:** For the type locality at Birness, Scotland.**Type Material:** n.d.**References:** (1) Jones, L.H.P. and A.A. Milne (1956) Birnessite, a new manganese oxide mineral from Aberdeenshire, Scotland. *Mineral. Mag.*, 31, 283–288. (2) (1957) *Amer. Mineral.*, 42, 440 (abs. ref. 1). (3) Glover, E.D. (1977) Characterization of a marine birnessite. *Amer. Mineral.*, 62, 278–285. (4) Chukhrov, F.V., A.I. Gorshkov, E.S. Rudnitskaya, V.V. Beresovskaya, and A.V. Sivtsov (1980) Manganese minerals in clay: a review. *Clays and Clay Minerals*, 28, 346–453. (5) Post, J.E. and D.R. Veblen (1990) Crystal structure determinations of synthetic sodium, magnesium, and potassium birnessite using TEM and the Rietveld method. *Amer. Mineral.*, 75, 477–489.

All rights reserved. No part of this publication may be reproduced, stored in a retrieval system or transmitted in any form or by any means, electronic, mechanical, photocopying, recording, or otherwise without the prior written permission of Mineral Data Publishing.