


Chapmanite



©2001 Mineral Data Publishing, version 1.2

Crystal Data: Monoclinic. *Point Group:* m . Crystals lath-shaped; powdery, compact, massive.

Physical Properties: *Fracture:* Conchoidal to irregular when compact. Hardness = 2.5 when compact. $D(\text{meas.}) = 3.69\text{--}3.75$ $D(\text{calc.}) = [4.29]$

Optical Properties: Semitransparent. *Color:* Olive-green to deep yellow. *Streak:* Greenish yellow to deep yellow. *Luster:* Nonmetallic.

Optical Class: Biaxial. $\alpha = 1.85 \perp$ flat face. $\beta = \text{n.d.}$ $\gamma = 1.96 \parallel$ length. $2V(\text{meas.}) = \text{n.d.}$

Cell Data: *Space Group:* Cm . $a = 5.19$ $b = 8.99$ $c = 7.70$ $\beta = 100^\circ 40'$ $Z = 2$

X-ray Powder Pattern: Keeley mine, Canada; could be confused with bismutoferrite. 7.63 (100), 3.58 (100), 3.88 (90), 3.19 (90), 2.90 (70), 2.59 (70), 2.54 (35)

Chemistry:

	(1)	(2)	(3)
SiO ₂	27.63	26.34	27.65
Al ₂ O ₃	0.22	0.50	
Fe ₂ O ₃	38.98	38.13	36.74
Sb ₂ O ₃	31.04	30.53	33.54
H ₂ O ⁺	1.89	3.62	2.07
H ₂ O [−]	0.31	0.64	
Total	100.07	99.76	100.00

(1) Smilkov, Czech Republic; corresponding to $(\text{Sb}_{0.93}^{3+}\text{Fe}_{0.07}^{3+})_{\Sigma=1.00}\text{Fe}_{2.00}^{2+}(\text{Si}_{1.98}\text{Al}_{0.02})_{\Sigma=2.00}\text{O}_{8.00}(\text{OH})$. (2) Do. (3) $\text{SbFe}_2(\text{SiO}_4)_2(\text{OH})$.

Occurrence: In veinlets cutting gneisses containing varying amounts of graphite (Smilkov, Czech Republic).

Association: Silver (Keeley mine, Canada).

Distribution: In the Keeley mine, South Lorrain Township, Ontario, Canada. In Mexico, at Velardeña, and in the La Papa mine, Victoria, Hidalgo, Durango. At Smilkov, near Votice, Czech Republic. In Germany, at Bräunsdorf, Freiberg, and Schneeberg, Saxony. From near Tafone, Tuscany, Italy. In the Mine de la Bessade, Massiac, Haute-Loire, France. At the Suzuyama mine, Kagoshima Prefecture, Japan.

Name: To honor Edward John Chapman (1821–1904), Professor of Geology, University of Toronto, Toronto, Canada.

Type Material: Royal Ontario Museum, Toronto, Canada, M14075; Harvard University, Cambridge, Massachusetts, USA, 91501.

References: (1) Walker, T.L. (1924) Chapmanite, a new mineral from South Lorraine, Ontario. Univ. of Toronto, Geol. Studies, 17, 5–8. (2) Walker, T.L. (1924) Chapmanite, a new hydrous ferrous silicoantimonate, from South Lorraine, Ontario. Contr. to Canadian Mineralogy, 5. (3) (1925) Amer. Mineral., 10, 40 (abs. ref. 2) (4) Milton, C., J.M. Axelrod, and B. Ingram (1958) Bismutoferrite, chapmanite, and “hypochlorite”. Amer. Mineral., 43, 656–670. (5) Čech, F. and P. Povondra (1963) Study of chapmanite from Smilkov, near Votice (Bohemia, Czechoslovakia). Acta Univ. Carolinae, Geol., 2, 97–114 (in Czech with English abs.). (6) (1964) Amer. Mineral., 49, 1499 (abs. ref. 5). (7) Zhukhlistov, A.P. and B.B. Zvyagin (1977) Determination of the crystal structures of chapmanite and bismuthoferrite [bismutoferrite] by high-voltage electron diffraction. Kristallografiya (Sov. Phys. Crystal.), 22, 731–738 (in Russian).

All rights reserved. No part of this publication may be reproduced, stored in a retrieval system or transmitted in any form or by any means, electronic, mechanical, photocopying, recording, or otherwise without the prior written permission of Mineral Data Publishing.