

Crystal Data: Orthorhombic. *Point Group:* $mm2$. As lamellar inclusions in löllingite, to 1.4 mm, or intergrown with paracostibite.

Physical Properties: Hardness = n.d. VHN = 781 (15 g load). $D(\text{meas.}) = \text{n.d.}$
 $D(\text{calc.}) = 6.89$

Optical Properties: Opaque. *Color:* In polished section, grayish. *Luster:* Metallic.
Pleochroism: Weak, pale gray with bluish to brownish tints. *Anisotropism:* Weak, reddish brown or orange, to bluish.

R_1 – R_2 : n.d.

Cell Data: *Space Group:* $Pmn2_1$. $a = 3.603$ $b = 4.868$ $c = 5.838$ $Z = 2$

X-ray Powder Pattern: Broken Hill, Australia.
 2.596 (100), 2.503 (90), 1.908 (80), 2.902 (60), 4.86 (50), 1.803 (50), 3.08 (40)

Chemistry:	(1)	(2)	(3)
Co	26.7	25.6	27.70
Fe	0.6	0.8	
Ni	0.2	2.4	
Sb	57.0	56.8	57.23
As	0.3	0.3	
S	15.1	14.7	15.07
Total	99.9	100.6	100.00

(1) Consols mine, Australia; by electron microprobe, average of three grains, corresponding to $(\text{Co}_{0.96}\text{Fe}_{0.02}\text{Ni}_{0.01})_{\Sigma=0.99}(\text{Sb}_{0.99}\text{As}_{0.01})_{\Sigma=1.00}\text{S}_{1.00}$. (2) Getön deposit, Sweden; by electron microprobe; corresponding to $(\text{Co}_{0.95}\text{Ni}_{0.09}\text{Fe}_{0.03})_{\Sigma=1.07}(\text{Sb}_{1.02}\text{As}_{0.01})_{\Sigma=1.03}\text{S}_{1.00}$. (3) CoSbS.

Polymorphism & Series: Dimorphous with paracostibite.

Mineral Group: Löllingite group.

Occurrence: Intimately intergrown with other hydrothermal sulfides (Broken Hill, Australia); in Pb–Zn–Cu–Ag ore deposits remobilized by hydrothermal solutions from later granite emplacement (Bergslagen, Sweden).

Association: Löllingite, willyamite, dyscrasite, ullmannite, pyrrargyrite (Broken Hill, Australia); nisbite, paracostibite, chalcopyrite, bismuth, pyrrhotite, galena, sphalerite, gersdorffite, ullmannite (Bergslagen, Sweden).

Distribution: In Australia, at the Consols mine, Broken Hill, New South Wales [TL]. From the Gruvåsen and Getön deposits, Bergslagen metallic province, Sweden. At Sulitjelma, northern Norway.

Name: For CObalt and *antimony*, STIBium, in its composition.

Type Material: National Museum of Natural History, Washington, D.C., USA, R849A.

References: (1) Cabri, L.J., D.C. Harris, and J.M. Stewart (1970) Costibite (CoSbS), a new mineral from Broken Hill, N.S.W., Australia. *Amer. Mineral.*, 55, 10–17. (2) Rowland, J.F., E.J. Gabe, and S.R. Hall (1975) The crystal structures of costibite (CoSbS) and paracostibite (CoSbS). *Can. Mineral.*, 13, 188–196. (3) Zakrzewski, M.A., E.A.J. Burke, and H.W. Nugteren (1980) Cobalt minerals in the Hallëfors area, Bergslagen, Sweden: new occurrences of costibite, paracostibite, nisbite and cobaltian ullmannite. *Can. Mineral.*, 18, 165–171.

All rights reserved. No part of this publication may be reproduced, stored in a retrieval system or transmitted in any form or by any means, electronic, mechanical, photocopying, recording, or otherwise without the prior written permission of Mineral Data Publishing.