(c)2001-2005 Mineral Data Publishing, version 1

Crystal Data: Triclinic. Point Group:  $\overline{1}$  (probable) or 1. As rosettes of crystals, to 0.6 mm, tabular on  $\{010\}$ , showing forms  $\{101\}$ ,  $\{10\overline{1}\}$ , and less commonly  $\{100\}$  and  $\{001\}$ , rectangular or square in outline; also finely granular and as porcelaneous coatings.

**Physical Properties:** Cleavage:  $\{010\}$ , perfect;  $\{100\}$ , fair. Tenacity: Flexible. Hardness = 2.5-3 D(meas.) = 2.48(1) D(calc.) = 2.482

**Optical Properties:** Transparent to opaque. Color: Colorless, white, creamy, bluish white. Streak: White. Luster: Vitreous to pearly if colorless; otherwise dull. Optical Class: Biaxial (+). Orientation: X (90°,41°); Y (240°,53°); Z (343°,74°) [with c (0°,0°) and  $b^*$  (0°,90°) using  $(\phi,\rho)$ ]. Dispersion: r > v, moderately strong.  $\alpha = 1.545(1)$   $\beta = 1.553(1)$   $\gamma = 1.566(1)$  2V(meas.) = 77° 2V(calc.) = 76.8°

**Cell Data:** Space Group:  $P\overline{1}$  (probable), or P1. a = 5.002(1) b = 5.175(1) c = 4.980(2)  $\alpha = 97.50(1)^{\circ}$   $\beta = 118.60(1)^{\circ}$   $\gamma = 104.74(1)^{\circ}$  Z = 2

**X-ray Powder Pattern:** Mont Saint-Hilaire, Canada; minor preferred orientation. 4.794 (100), 2.360 (40), 1.972 (30), 1.857 (30), 1.842 (30), 4.296 (20), 4.182 (20)

$\alpha$	•	
Chei	nic	trw
CHO	.1113	υт.у •

	(1)	(2)	(3)	(4)
$SiO_2$		0.03	3.22	
$Al_2O_3$	65.2	63.7	59.6	65.36
FeO		0.14	0.08	
MgO			0.96	
CaO	0.48		0.23	
$Na_2O$		0.09	0.35	
$\mathrm{H_2O}$	35.76	35.76	[35.56]	34.64
Total	101.44	99.72	[100.00]	100.00

(1) Mont Saint-Hilaire, Canada;  $H_2O$  by TGA,  $(OH)^{1-}$  confirmed by IR. (2) Do.; by electron microprobe,  $H_2O$  from (1). (3) Francon quarry, Canada; by electron microprobe, average of four analyses,  $H_2O$  by difference. (4) Al $(OH)_3$ .

Polymorphism & Series: Polymorphous with bayerite, gibbsite, and nordstrandite.

Occurrence: A late-stage hydrothermal mineral lining vugs in veins in nepheline syenite associated with an intrusive alkalic gabbro-syenite complex (Mont Saint-Hilaire, Canada); in vugs within silicocarbonatite sills intruding limestones (Francon quarry, Canada); in miarolitic cavities in a nepheline syenite sill (near Saint-Amable, Canada).

Association: Calcite, albite, pyrite (Mont Saint-Hilaire, Canada); weloganite, calcite, quartz, albite, pyrite, cryolite, strontianite, dresserite, dawsonite, fluorite, analcime (Francon quarry, Canada); microcline, eudialyte, yofortierite, astrophyllite, serandite, aegirine (near Saint-Amable, Canada).

**Distribution:** Found at Mont Saint-Hilaire, in the Francon quarry, Montreal Island, Montreal, and near Saint-Amable, Quebec, Canada.

Name: Honors Canadian physician E.J. Doyle, of Ottawa, Canada, who found the mineral at Mont Saint-Hilaire, Canada.

**Type Material:** Canadian Museum of Nature, Ottawa, 48932; Canadian Geological Survey, Ottawa; Royal Ontario Museum, Toronto, Canada, M41025; National Museum of Natural History, Washington, D.C., USA, 162728.

**References:** (1) Chao, G.Y., J. Baker, A.P. Sabina, and A.C. Roberts (1985) Doyleite, a new polymorph of Al(OH)<sub>3</sub>, and its relationship to bayerite, gibbsite and nordstrandite. Can. Mineral., 23, 21–28. (2) (1986) Amer. Mineral., 71, 845 (abs. ref. 1).

All rights reserved. No part of this publication may be reproduced, stored in a retrieval system or transmitted in any form or by any means, electronic, mechanical, photocopying, recording, or otherwise without the prior written permission of Mineral Data Publishing.