

**Crystal Data:** Monoclinic. *Point Group:* 2/m, 2 or m. Crystals flattened on [001] mostly split and curved, prismatic to elongated lamellar, or curved ribbon-like, to 3.2 mm; striations on {001} across elongation; as fan- and rosette-like clusters, or chaotic aggregates to 6.5 mm.

**Physical Properties:** *Cleavage:* Perfect on {001}. *Fracture:* n.d. *Tenacity:* Flexible, inelastic. Hardness = < 1 VHN = 13 (2 g load). D(meas.) = n.d. D(calc.) = 3.467

**Optical Properties:** Opaque. *Color:* Dark bronze (fresh), to nearly black; in reflected light, gray with a bluish to pale beige tint. *Streak:* Black. *Luster:* Metallic (fresh), dull, or tarnishes to iridescent purplish or golden-brown.

*Optical Class:* n.d. *Birefractance:* Distinct. *Anisotropism:* Distinct, gray-bluish to yellowish beige.

R<sub>1</sub>-R<sub>2</sub>: (470) 11.4-11.6, (546) 11.2-12.4, (589) 11.1-13.6, (650) 11.0-15.5

**Cell Data:** *Space Group:* C2/m, Cm, or C2. *a* = 5.463(5) *b* = 15.865(17) *c* = 10.825(12)  $\beta$  = 93.7(1)° *Z* = 2

**X-ray Powder Pattern:** Oktyabr'skiy mine, Norilsk district, Krasnoyarskiy Kray, Russia. 5.392 (100), 10.83 (13), 2.696 (12), 2.524 (12), 1.837 (11), 2.152 (8), 3.281 (7)

<b>Chemistry:</b>	(1)
Mg	0.02
Fe	61.92
Ni	0.03
Cu	0.09
S	19.45
O	16.3
H	[1.03]
Total	98.84

(1) Oktyabr'skiy mine, Norilsk district, Krasnoyarskiy Kray, Russia; average of 9 electron microprobe analyses, Fe<sup>2+</sup>/Fe<sup>3+</sup> calculated for charge balance, H calculated as if present only as OH, presence of OH and absence of H<sub>2</sub>O confirmed by IR spectroscopy; corresponding to (Fe<sub>5.98</sub>Cu<sub>0.015</sub>Ni<sub>0.005</sub>)<sub>Σ=6.00</sub>S<sub>6</sub>(Fe<sub>4.89</sub>Mg<sub>0.01</sub>)<sub>Σ=4.90</sub>(OH)<sub>9.80</sub>Fe<sup>3+</sup><sub>0.09</sub>(OH)<sub>0.27</sub>.

**Occurrence:** Of low-temperature hydrothermal origin coating cavities in pentlandite-mooihoekite-cubanite ore with minor magnetite and chalcopyrite.

**Association:** Ferrovalleriite, magnetite, an Fe-rich chlorite-type phyllosilicate.

**Distribution:** From Shaft no 1, Oktyabr'skiy mine, Oktyabr'skoye Cu-Ni-PGM deposit, Talnakh, Norilsk district, Krasnoyarskiy Kray, Siberia, Russia.

**Name:** As the structural analogue (based on chemical, X-ray, and IR data similarities) of *tochilinite* with essential *ferrous iron*.

**Type Material:** A.E. Fersman Mineralogical Museum, Academy of Sciences, Moscow, Russia (4058/1).

**References:** (1) Pekov, I.V., E.V. Sereda, Yu.S. Polekhovsky, S.N. Britvin, N.V. Chukanov, V.O. Yapaskurt, and I.A. Bryzgalov (2012) Ferrotochilinite, 6FeS·5Fe(OH)<sub>2</sub>, a new mineral from Oktyabr'skoye ore deposit (Norilsk ore district, Siberia, Russia). Zap. Ross. Mineral. Obsch., 141(4), 1-11 (in Russian, with English abstract). (2) (2014) Amer. Mineral., 99, 242-243 (abs. ref. 1).