

**Crystal Data:** Orthorhombic. *Point Group:* mm2. As 100-250  $\mu\text{m}$  lamellar crystals filling interstitial areas within gehlenite or intergrown with rankinite and Ti-rich andradite.

**Physical Properties:** *Cleavage:* None. *Fracture:* Irregular. *Tenacity:* Brittle. *Hardness* = 5-5.5 VHN = 706 (50 g load). D(meas.) = n.d. D(calc.) = 3.264

**Optical Properties:** Transparent. *Color:* Pale yellow or gray. *Streak:* White. *Luster:* Vitreous. *Optical Class:* Uniaxial (+).  $\omega = 1.634(2)$   $\varepsilon = 1.640(2)$  *Orientation:*  $Z = E$ .

**Cell Data:** *Space Group:*  $Pnm2_1$ .  $a = 9.3845(6)$   $b = 21.7310(14)$   $c = 6.8346(4)$   $Z = 4$

**X-ray Powder Pattern:** Calculated pattern.

2.713 (100), 2.765 (44), 2.759 (42), 1.762 (32), 2.518 (29), 2.402 (23), 2.897 (19)

<b>Chemistry:</b>	(1)
CaO	59.76
SiO <sub>2</sub>	28.87
Al <sub>2</sub> O <sub>3</sub>	0.04
FeO	0.15
MgO	0.16
BaO	0.05
SrO	0.24
V <sub>2</sub> O <sub>5</sub>	0.10
P <sub>2</sub> O <sub>5</sub>	7.38
Na <sub>2</sub> O	1.55
K <sub>2</sub> O	1.73
Total	100.03

(1) Hatrurim Basin, Negev Desert, Israel; average of 21 electron microprobe analyses supplemented by Raman spectroscopy; corresponds to  $[\text{Ca}_{1.82}\text{Na}_{0.09}\text{K}_{0.06}(\text{Mg},\text{Fe},\text{Sr},\text{Ba})_{0.02}]_{\Sigma=1.99}(\text{Si}_{0.82}\text{P}_{0.18})_{\Sigma=1.00}\text{O}_4$ .

**Occurrence:** In Ca- and Al-rich paralava, an ultrahigh-temperature combustion metamorphic rock.

**Association:** Larnite (partially hydrated), gehlenite, rankinite, Ti-rich andradite, ferrian perovskite, magnesioferrite, hematite, ettringite.

**Distribution:** From the southern Hatrurim Basin, Negev Desert, Israel.

**Name:** Derived from “flame”, in allusion to the origin of the mineral by ultrahigh-temperature metamorphism from the natural combustion of fossil fuel.

**Type Material:** Central Siberian Geological Museum, V.S. Sobolev Institute of Geology and Mineralogy, Novosibirsk, Russia (XIII-341/1).

**References:** (1) Sokol, E.V., Y.V. Seryotkin, S.N. Kokh, Y. Vapnik, E.N. Nigmatulina, S.V. Goryainov, E.V. Belogub, and V.V. Sharygin (2015) Flamite,  $(\text{Ca},\text{Na},\text{K})_2(\text{Si},\text{P})\text{O}_4$ , a new mineral from ultrahigh-temperature combustion metamorphic rocks, Hatrurim Basin, Negev Desert, Israel. *Mineral. Mag.*, 79(3), 583-596. (2) Gfeller, F., R. Widmer, B. Krüger, E.V. Galuskin, I.O. Galuskina, and T. Armbruster (2015) The crystal structure of flamite and its relation to  $\text{Ca}_2\text{SiO}_4$  polymorphs and nagelschmidite. *Eur. J. Mineral.*, 27, 755-769. (3) (2016) Amer. *Mineral.*, 101, 1713 (abs. refs. 1 & 2).