©2001 Mineral Data Publishing, version 1.2

Crystal Data: Triclinic. Point Group: $\overline{1}$. Crystals, < 1 mm, slender needles and plates, elongated along [001] with $\{010\}$, $\{100\}$, and $\{\overline{1}10\}$. Twinning: On $\{100\}$.

Cleavage: {010}, rather perfect; {100}, faint. Fracture: Brittle. **Physical Properties:** Hardness = 5 D(meas.) = 3.148 D(calc.) = 3.157

Optical Properties: Semitransparent. Color: Wine-yellow, honey-yellow, colorless; light wine-yellow in thin section. Luster: Vitreous to adamantine. Optical Class: Biaxial (+). Pleochroism: X = colorless; Z = wine-yellow. Dispersion: r > v, Absorption: Z > Y > X. $n = \sim 1.7$, birefringence ~ 0.012 . 2V(meas.) = Large.strong.

Cell Data: Space Group: $P\overline{1}$. a = 5.676(2) b = 7.259(3) c = 9.586(3) $\alpha = 101.08(4)^{\circ}$ $\beta = 101.14(4)^{\circ}$ $\gamma = 90.27(4)^{\circ}$ Z = 1

X-ray Powder Pattern: Hradiste Mountain, Czech Republic. 2.961(100), 3.073(90), 1.896(70), 3.961(50), 2.628(50), 2.493(50), 1.822(50)

	(1)		(1)
SiO_2	32.07	Nb_2O_5	1.38
TiO_2	8.06	Ta_2O_5	0.22
$ m ZrO_2$	6.58	MnO	2.06
Al_2O_3	0.03	CaO	32.05
La_2O_3	0.60	Na_2O	7.46
Ce_2O_3	1.01	\mathbf{F}	12.09
Fe_2O_3	1.25	$-O = F_2$	5.09
		Total	99.77

(1) Hradiste Mountain, Czech Republic; by electron microprobe, average of fourteen analyses; corresponds to $(Na_{3.52}Ca_{0.48})_{\Sigma=4.00}Ca_{8.00}RE_{0.15}(Ti_{1.52}Zr_{0.80}Mn_{0.45}Fe_{0.25}Nb_{0.09}Ta_{0.02}Al_{0.01})_{\Sigma=3.14}$ $Si_{7.80}[O_{27.68}F_{8.32}]_{\Sigma=36.00}.$

Occurrence: In cavities and embedded in the groundmass of a phonolite.

Association: Aegirine.

Distribution: From Hradiste Mountain, near Mildenau, Czech Republic.

Name: For the locality, formerly named Hoher Hain, in the Czech Republic.

Type Material: Charles University, Prague, Czech Republic, 20530.

References: (1) Dana, E.S. (1899) Dana's system of mineralogy, (6th edition), app. I, 31–32. (2) Vlasov, K.A., Ed. (1966) Mineralogy of rare elements, v. II, 385. (3) Johan, Z. and F. Cech (1989) New data on hainite, $Na_2Ca_4[(Ti, Zr, Mn, Fe, Nb, Ta)_{1.50} \square_{0.50}](Si_2O_7)_2F_4$ and its crystallochemical relationship with götzenite, $Na_2Ca_5Ti(Si_2O_7)_2F_4$. Compt. Rendus Acad. Sci. Paris, 308, 1237–1242 (in French with English abs.). (4) (1990) Amer. Mineral., 75, 936 (abs. ref. 3).