


Manganbabingtonite



©2001 Mineral Data Publishing, version 1.2

Crystal Data: Triclinic. *Point Group:* $\bar{1}$. Crystals short prismatic, to 5 mm.

Physical Properties: *Cleavage:* Perfect on {001}, less perfect on {010}, poor on {100}. *Fracture:* Subconchoidal. *Tenacity:* Brittle. Hardness = 6.5–7 D(meas.) = 3.452 D(calc.) = [3.59]

Optical Properties: Translucent to opaque. *Color:* Dark greenish black to yellowish green. *Optical Class:* Biaxial (+). *Pleochroism:* X = emerald-green; Y = faint rose to nearly colorless; Z = purple-brown to rose-brown. *Orientation:* Y = \simeq c; Z = b. *Dispersion:* r > v, strong. $\alpha = 1.716$ $\beta = 1.730$ $\gamma = 1.746$ 2V(meas.) = 78°–82° 2V(calc.) = 86°

Cell Data: *Space Group:* $P\bar{1}$. a = 6.88 b = 11.80 c = 6.77 $\alpha = 90^\circ 30'$ $\beta = 93^\circ 30'$ $\gamma = 104^\circ 54'$ Z = 2

X-ray Powder Pattern: Rudnyi Kaskad deposit, Russia.

3.104 (10), 2.998 (10), 2.938 (10), 2.162 (10), 3.442 (8), 1.654 (7), 1.427 (6)

Chemistry:	(1)	(2)	(1)	(2)
SiO ₂	51.85	52.45	K ₂ O	0.09
TiO ₂	0.15		F	0.12
Al ₂ O ₃	0.56		Cl	0.06
Fe ₂ O ₃	12.26	13.94	H ₂ O ⁺	1.95
FeO	4.52	6.27	H ₂ O ⁻	0.26
MnO	7.91	6.19	H ₂ O	1.57
MgO	0.60		CO ₂	0.13
CaO	19.00	19.58	–O = (F, Cl) ₂	0.06
Na ₂ O	0.25			
			Total	99.65 100.00

(1) Rudnyi Kaskad deposit, Russia. (2) Ca₂(Mn²⁺, Fe²⁺)Fe³⁺Si₅O₁₄(OH), calculated for Mn²⁺:Fe²⁺ = 1:1.

Polymorphism & Series: Forms a series with babingtonite.

Occurrence: In thin stringers in amphibolized garnet-pyroxene-magnetite skarn associated with an iron deposit (Rudnyi Kaskad deposit, Russia); replacing hedenbergite-johannsenite skarn (Iron Cap mine, Arizona, USA).

Association: Epidote, calcite, quartz (Rudnyi Kaskad deposit, Russia); ilvaite, andradite, epidote, thaumasite, calcite, quartz, pyrite, galena, sphalerite (Iron Cap mine, Arizona, USA).

Distribution: From the Rudnyi Kaskad deposit, eastern Sayan Mountains, Siberia, Russia. In the Iron Cap Pb–Zn mine, near Aravaipa, Graham Co., Arizona, USA.

Name: For the manganese content and relation to *babingtonite*.

Type Material: A.E. Fersman Mineralogical Museum, Academy of Sciences, Moscow, Russia, 72039.

References: (1) Vinogradova, R.A., V.A. Sychkova, and Y.K. Kabalov (1966) Manganiferous babingtonite from the Rudnyi Kaskad deposit, Eastern Sayan. Doklady Acad. Nauk SSSR, 169, 434–437 (in Russian). (2) Vinogradova, R.A. and I.N. Plyusina (1967) Composition, properties, and crystallochemical features of minerals of the isomorphous series ferrobabingtonite-manganbabingtonite. Vestn. Moskov. Univ. Ser. IV, Geol., 54–67 (in Russian). (3) (1968) Amer. Mineral., 53, 1064–1065 (abs. refs. 1 and 2). (4) Burt, D.M. and D. London (1978) Manganbabingtonite and ilvaite from a hedenbergite-johannsenite skarn, Aravaipa, Arizona. Geol. Soc. Amer., Abs. with Prog., 10, 98 (abs.).

All rights reserved. No part of this publication may be reproduced, stored in a retrieval system or transmitted in any form or by any means, electronic, mechanical, photocopying, recording, or otherwise without the prior written permission of Mineral Data Publishing.