©2001 Mineral Data Publishing, version 1.2

**Crystal Data:** Monoclinic. *Point Group:* n.d. As matted laths and fibers, elongated along [010], forming small rosettes, to 2 mm.

**Physical Properties:** Cleavage:  $\{001\}$  (?). Hardness = 3 D(meas.) = 2.36 D(calc.) = 2.47

**Optical Properties:** Transparent to translucent. *Color:* White. *Luster:* Silky. *Optical Class:* Biaxial (+) (probable). *Orientation:*  $Y = b; Z \wedge c = 18^{\circ}.$   $\alpha = 1.500-1.504$   $\beta = 1.505-1.510$   $\gamma = 1.513-1.519$  2V(meas.) = Moderate to large.  $2V(\text{calc.}) = 76^{\circ}$ 

Cell Data: Space Group: n.d. a=13.51 b=6.55 c=13.51  $\beta=104^{\circ}$  Z=8

**X-ray Powder Pattern:** Kimberley, South Africa. 2.94 (vvs), 6.6 (vs), 13.1 (s), 4.67 (s), 2.80 (ms), 1.967 (ms), 4.18 (m)

Chemistry:

	(1)	(2)
$\mathrm{SiO}_2$	58.5	51.57
${ m Al}_2{ m O}_3$	0.0	2.31
CaO	13.4	10.07
MgO	0.2	0.00
$Na_2O$	7.9	10.15
$K_2O$	6.0	4.43
$\overline{\mathrm{H_2O^+}}$		9.74
$H_2^-O^-$		8.02
$\overline{\text{H}_2\text{O}}$	13.4	
$\overline{\text{CO}}_2$		0.87
LOĨ		2.47
Total	99.4	99.63

(1) Kimberley, South Africa; corresponds to  $(Na_{1.04}Ca_{0.99}K_{0.54}Mg_{0.02})_{\Sigma=2.59}$   $Si_{4.02}O_{9.99} \bullet 3.09H_2O.$  (2) Lovozero massif, Russia; corresponds to  $(Na_{1.28}Ca_{0.80}K_{0.42})_{\Sigma=2.50}$   $(Si_{3.80}Al_{0.20})_{\Sigma=4.00}O_{9.99} \bullet 2.75H_2O.$ 

Occurrence: In a kimberlite pipe (Kimberley, South Africa); in a differentiated alkalic massif (Lovozero massif, Russia).

Association: Rhodesite (Kimberley, South Africa); natrolite, steenstrupine, chkalovite, neptunite, sérandite, aegirine, leucosphenite, zorite, raite, polylithionite, halite (Lovozero Massif, Russia).

**Distribution:** In the Bultfontein diamond mine, Kimberley, Cape Province, South Africa. On Mt. Karnasurt, Lovozero massif, Kola Peninsula, Russia. In the Höwenegg quarry, Hegau, Baden-Württemberg, Germany.

Name: For Edgar Donald Mountain, Professor of Geology, Rhodes University, Grahamstown, South Africa.

**Type Material:** The Natural History Museum, London, England, 1957,369; National Museum of Natural History, Washington, D.C., USA, 114792.

References: (1) Gard, J.A., H.F.W. Taylor, and R.A. Chalmers (1957) An investigation of two new minerals: rhodesite and mountainite. Mineral. Mag., 31, 611–623. (2) (1958) Amer. Mineral., 43, 624–625 (abs. ref. 1). (3) Bussen, I.V., L.G. Latysheva, Y.P. Men'shikov, A.N. Mer'kov, T.S. Romanova, and A.S. Sakharov (1973) First find of mountainite in the Soviet Union. Doklady Acad. Nauk SSSR, 210, 674–677 (in Russian).

All rights reserved. No part of this publication may be reproduced, stored in a retrieval system or transmitted in any form or by any means, electronic, mechanical, photocopying, recording, or otherwise without the prior written permission of Mineral Data Publishing.