

Pickeringite

MgAl₂(SO₄)₄•22H₂O

©2001-2005 Mineral Data Publishing, version 1

Crystal Data: Monoclinic. *Point Group:* 2. Acicular to hairlike crystals, with many forms measured although terminated crystals are very rare; in radial or matted aggregates; typically as incrustations and efflorescences.

Physical Properties: *Cleavage:* Poor on {010}. *Fracture:* Conchoidal. *Tenacity:* Brittle. Hardness = 1.5 D(meas.) = 1.73–1.79 D(calc.) = 1.84 Soluble in H₂O, astringent taste.

Optical Properties: Semitransparent. *Color:* Colorless, white; may be pale shades of yellow, green, or red from metallic impurities; colorless in transmitted light. *Luster:* Vitreous. *Optical Class:* Biaxial (-). *Orientation:* Y = b; Z ∧ a = 36°. α = 1.475 β = 1.480 γ = 1.483 2V(meas.) = 60°

Cell Data: *Space Group:* P2₁/c. a = 6.1844(2) b = 24.2715(9) c = 21.2265(7) β = 100.326(4)° Z = 4

X-ray Powder Pattern: Tucumcari, New Mexico, USA. (ICDD 12-299). 4.82 (100), 3.510 (90), 4.32 (35), 4.122 (30), 3.791 (30), 6.08 (20), 4.97 (20)

Chemistry:	(1)	(2)
SO ₃	37.84	37.29
Al ₂ O ₃	12.30	11.87
MgO	4.35	4.69
CaO	0.09	
H ₂ O	44.66	46.15
insol.	0.50	
Total	99.74	100.00

(1) Quetena, Chile. (2) MgAl₂(SO₄)₄•22H₂O.

Polymorphism & Series: Forms a series with halotrichite.

Mineral Group: Halotrichite group.

Occurrence: A common secondary mineral formed by alteration of pyrite in aluminous rocks or in coal seams; in the oxidized zone of pyritic hydrothermal mineral deposits, typically in arid regions, typically post-mining; a fumarolic product; formed in caves.

Association: Kalinite, alunogen, epsomite, melanterite, copiapite, gypsum.

Distribution: Widespread, so only a few localities are listed. In Chile, abundant from Cerros Pintados, 80 km southeast of Iquique, Tarapacá; at Quetena, west of Calama, and Chuquicamata, Antofagasta. In the USA, in New Mexico, from near Tucumcari, Quay Co.; at The Geysers, Sonoma Co., California; from Alum Point, Salt Lake Co., Utah. In Canada, at Newport, Nova Scotia, and from the junction of the two main branches of the Smoky River, Alberta. In Germany, at Wetzelsstein, near Saalfeld, and from near Lehesten, Thuringia. On Valachov Hill, near Skřivava, Czech Republic. At Cervencia (Opáľbánya), Slovakia. In Italy, from Baia di Levante, Vulcano, Lipari Islands; on Mt. Etna, Sicily; and on Elba. At volcanoes on the Kamchatka Peninsula, Russia.

Name: To honor John Pickering (1777–1846), American lawyer and philologist of Boston, Massachusetts, USA.

References: (1) Palache, C., H. Berman, and C. Frondel (1951) Dana's system of mineralogy, (7th edition), v. II, 523–526. (2) Quartieri, S., M. Triscari, and A. Viani (2000) Crystal structure of the hydrated sulphate pickeringite MgAl₂(SO₄)₄•22H₂O: X-ray powder diffraction study. Eur. J. Mineral., 12, 1131–1138.

All rights reserved. No part of this publication may be reproduced, stored in a retrieval system or transmitted in any form or by any means, electronic, mechanical, photocopying, recording, or otherwise without the prior written permission of Mineral Data Publishing.