©2001 Mineral Data Publishing, version 1.2

Crystal Data: Monoclinic. Point Group: 2/m. As dense, compact masses of tiny lathlike crystals, which may be in parallel growth; as platy aggregates.

Physical Properties: Cleavage: Perfect on $\{001\}$. Hardness = <3 D(meas.) = 3.433 D(calc.) = [3.44]

Optical Properties: Translucent. Color: White to pale pink; colorless in thin section.

Luster: Dull to somewhat resinous.

Optical Class: Biaxial (+). Dispersion: r < v, slight. $\alpha = 1.654(1)$ $\beta = 1.660(1)$ $\gamma = 1.678(1)$ $2V(meas.) = 61(2)^{\circ}$ $2V(calc.) = 61^{\circ}$

Cell Data: Space Group: C2/m. a = 13.208(4) b = 8.287(2) c = 13.089(9) $\beta = 106.65(6)^{\circ}$ Z = 2

X-ray Powder Pattern: Långban, Sweden. (ICDD 16-411). 3.15 (100), 4.20 (80), 3.04 (55), 2.102 (40), 6.47 (35), 6.29 (35), 2.947 (35)

\sim	i	. ,	
	hem	ıst	rv:

	(1)	(2)	(3)	(4)
SiO_2	23.58	23.57	24.6	24.62
$Mn\bar{O}$	2.48	2.49	4.4	4.84
PbO	31.03	30.04	30.7	30.48
CaO	25.95	23.12	22.6	22.98
$\operatorname{Sr} \operatorname{O}$	1.40	2.79	0.7	
Na_2O	0.40			
K_2O	0.13			
$\overline{\mathrm{H}_{2}^{-}\mathrm{O}^{+}}$	6.35	6.15	[6.13]	6.15
$\overline{\mathrm{H}_{2}^{-}\mathrm{O}^{-}}$		0.45		
$\overline{\mathrm{CO}_2}$		0.61		
SO_2	9.0			
SO_3		10.81	10.6	10.93
Total	100.32	100.03	99.73	100.00

(1–2) Franklin, New Jersey, USA. (3) Långban, Sweden; by electron microprobe, H_2O calculated from stoichiometry. (4) $Pb_2Ca_6MnSi_6O_{18}(SO_4)_2(OH)_2 \cdot 4H_2O$.

Occurrence: As nodular masses in calcium silicate lenses containing abundant garnet (Franklin, New Jersey, USA); as fracture fillings in a metamorphosed Fe–Mn orebody (Långban, Sweden).

Association: Garnet, titanite, zircon, phlogopite, axinite, willemite, datolite, barite, calcite, arsenopyrite, sphalerite, rhodonite, rhodochrosite.

Distribution: From Franklin, Sussex Co., New Jersey, USA. At Långban, Värmland, Sweden.

Name: For Colonel Washington Augustus Roebling (1837–1926), of Trenton, New Jersey, USA, distinguished American civil engineer, mineral collector, and benefactor of American geological sciences.

Type Material: National Museum of Natural History, Washington, D.C., USA, R8824, 124351; The Natural History Museum, London, England, 1925,69.

References: (1) Dana, E.S. (1899) Dana's system of mineralogy, (6th edition), app. I, 60.

- (2) Blix, R. (1931) The chemical composition of roeblingite. Amer. Mineral., 16, 455–469.
- (3) Moore, P.B. and J. Shen (1984) Roeblingite, $Pb_2Ca_6(SO_4)_2(OH)_2(H_2O)_4[Mn(Si_3O_9)_2]$: its crystal structure and comments on the lone pair effect. Amer. Mineral., 69, 1173–1179.
- (4) Dunn, P.J. (1985) The lead silicates from Franklin, New Jersey: occurrence and composition. Mineral. Mag., 49, 721–727. (5) Braithwaite, R.S.W. (1985) Roeblingite: a revised formula from infra-red and thermal analysis data. Mineral. Mag., 49, 756–758.

All rights reserved. No part of this publication may be reproduced, stored in a retrieval system or transmitted in any form or by any means, electronic, mechanical, photocopying, recording, or otherwise without the prior written permission of Mineral Data Publishing.