


Schreyerite



©2001-2005 Mineral Data Publishing, version 1

Crystal Data: Monoclinic. *Point Group:* n.d. As lamellae and grains, to 30 μm , exsolved within rutile crystals. *Twinning:* Polysynthetic, universal.

Physical Properties: Hardness = n.d. VHN = 1100–1200 D(meas.) = n.d. D(calc.) = [4.46]

Optical Properties: Opaque. *Color:* Reddish brown; gray in reflected light. *Optical Class:* Biaxial. $n = 2.7$ *Pleochroism:* Weak; yellow-brown to reddish brown. *Anisotropism:* Moderate. *Birefractance:* White to brownish. R: (546) 21

Cell Data: *Space Group:* n.d. $a = 7.06(1)$ $b = 5.024(3)$ $c = 18.84(3)$ $\beta = 119.68(7)^\circ$ Z = [4]

X-ray Powder Pattern: Lasamba Hill, Kenya. 2.737 (vs), 2.874 (s), 4.075 (m), 3.381 (m), 2.432 (w), 2.518 (vvw), 2.310 (vvw)

Chemistry:	(1)	(2)	(3)
TiO ₂	61.07	60.95	61.52
Al ₂ O ₃	0.43	0.30	
V ₂ O ₃	35.93	33.80	38.48
Cr ₂ O ₃	2.23	2.45	
FeO	0.10	1.25	
MnO		0.18	
MgO	0.11		
Total	99.87	98.93	100.00

(1) Lasamba Hill, Kenya; by electron microprobe, average of six analyses, total Fe as FeO; corresponds to $(\text{V}_{1.86}\text{Cr}_{0.12}\text{Al}_{0.02})_{\Sigma=2.00}\text{Ti}_3\text{O}_9$. (2) Sättra, Sweden; by electron microprobe, average of four analyses, total Fe as FeO; corresponds to $(\text{V}_{1.8}\text{Cr}_{0.2})_{\Sigma=2.0}\text{Ti}_3\text{O}_9$. (3) V₂Ti₃O₉.

Polymorphism & Series: Dimorphous with kyzylkumite.

Occurrence: In quartz-biotite-sillimanite-diopside-epidote gneiss in contact with quartzite, formed above 600 °C and 5 kbar (Lasamba Hill, Kenya); in quartzitic schists (Ol'khon Gate Straits, Lake Baikal, Russia).

Association: Vanadian rutile, kornerupine, kyanite, sillimanite, muscovite, apatite, tourmaline, graphite, pyrrhotite, chalcopyrite, pentlandite (Lasamba Hill, Kenya); rutile, eskolaite, karelianite, vuorelainenite, olkhonskite (Ol'khon Gate Straits, Lake Baikal, Russia).

Distribution: From six km southeast of Lasamba Hill, Kwale district, south of Voi, Kenya. At Komolo, Tanzania. In Sweden, from the Sättra mine, Doverstorp, Bergslagen metallic province. Found 4.5 km south of the Ol'khon Gate Strait, on the western shore of Lake Baikal, Siberia, Russia.

Name: To honor Dr. Werner Schreyer (1930–), Professor of Mineralogy, Ruhr University, Bochum, Germany.

Type Material: University of Heidelberg, Heidelberg, Germany; National Museum of Natural History, Washington, D.C., USA, 136230.

References: (1) Medenbach, O. and K. Schmetzer (1976) Schreyerit (V₂Ti₃O₉), ein neues Vanadium-Mineral aus Kenya. *Naturwiss.*, 63, 293–294 (in German). (2) (1977) *Amer. Mineral.*, 62, 395 (abs. ref. 1). (3) Medenbach, O. and K. Schmetzer (1978) Schreyerite, V₂Ti₃O₉, a new mineral. *Amer. Mineral.*, 63, 1182–1186. (4) Zakrzewski, M.A., E.A.J. Burke, and W.J. Lustenhouwer (1982) Vuorelainenite, a new spinel, and associated minerals from the Sättra (Doverstorp) pyrite deposit, central Sweden. *Can. Mineral.*, 20, 281–290. (5) Bernhardt, H.-J., K. Schmetzer, and O. Medenbach (1983) Berdesinskiite, V₂TiO₅, a new mineral from Kenya and additional data for schreyerite, V₂Ti₃O₉. *Neues Jahrb. Mineral., Monatsh.*, 110–118.

All rights reserved. No part of this publication may be reproduced, stored in a retrieval system or transmitted in any form or by any means, electronic, mechanical, photocopying, recording, or otherwise without the prior written permission of Mineral Data Publishing.