


# Sigloite



©2001-2005 Mineral Data Publishing, version 1

**Crystal Data:** Triclinic. *Point Group:*  $\bar{1}$ . Crystals are pseudomorphic after paravauxite crystals, distorted and curved, showing {010}, {001}, {100}, {110}, several other forms, to about 1 cm, in intergrowths.

**Physical Properties:** *Cleavage:* On {010}, perfect; on {001}, good. Hardness = 3  
D(meas.) = 2.35 D(calc.) = [2.40]

**Optical Properties:** Semitransparent. *Color:* Pale straw-yellow, pale brownish orange, light brown.

*Optical Class:* Biaxial (+). *Orientation:* X (90°, 32°); Y (-144°, 70°); Z (-46°, 66°) [using ( $\phi, \rho$ )].  
*Dispersion:*  $r < v$ , strong.  $\alpha = 1.563$   $\beta = 1.586(1)$   $\gamma = 1.619$   $2V(\text{meas.}) = 76^\circ$

**Cell Data:** *Space Group:*  $P\bar{1}$ .  $a = 5.190(2)$   $b = 10.419(4)$   $c = 7.033(3)$   $\alpha = 105.00(3)^\circ$   
 $\beta = 111.31(3)^\circ$   $\gamma = 70.87(3)^\circ$   $Z = 1$

**X-ray Powder Pattern:** Siglo XX mine, Bolivia.  
9.69 (10), 6.46 (9), 4.86 (9), 3.23 (7), 2.82 (6), 2.56 (5), 4.68 (4b)

## Chemistry:

|                                | (1)    | (2)    |
|--------------------------------|--------|--------|
| P <sub>2</sub> O <sub>5</sub>  | 27.47  | 29.77  |
| SiO <sub>2</sub>               | 0.11   |        |
| Al <sub>2</sub> O <sub>3</sub> | 21.09  | 21.38  |
| Fe <sub>2</sub> O <sub>3</sub> | 13.53  | 16.74  |
| FeO                            | 2.76   |        |
| MnO                            | 0.24   |        |
| MgO                            | 0.87   |        |
| Na <sub>2</sub> O              | 0.44   |        |
| K <sub>2</sub> O               | 0.26   |        |
| H <sub>2</sub> O               | 33.55  | 32.11  |
| Total                          | 100.32 | 100.00 |

(1) Siglo XX mine, Bolivia. (2) FeAl<sub>2</sub>(PO<sub>4</sub>)<sub>2</sub>(OH)<sub>3</sub> · 7H<sub>2</sub>O.

**Mineral Group:** Paravauxite group.

**Occurrence:** A rare late-stage secondary mineral, formed by oxidation of vauxite, in open fractures cutting cassiterite veins in a hydrothermal tin deposit.

**Association:** Wavellite, metavauxite, crandallite, childrenite, vauxite, quartz.

**Distribution:** Found in the Siglo XX mine, Llallagua, Potosí, Bolivia.

**Name:** For its occurrence in the Siglo XX mine, Llallagua, Bolivia.

**Type Material:** Harvard University, Cambridge, Massachusetts, 107443; National Museum of Natural History, Washington, D.C., USA, 115320.

**References:** (1) Hurlbut, C.S., Jr. and R. Honea (1962) Sigloite, a new mineral from Llallagua, Bolivia. *Amer. Mineral.*, 47, 1–8. (2) Hawthorne, F.C. (1988) Sigloite: the oxidation mechanism in [M<sub>2</sub><sup>3+</sup>(PO<sub>4</sub>)<sub>2</sub>(OH)<sub>2</sub>(H<sub>2</sub>O)<sub>2</sub>]<sup>2-</sup> structures. *Mineral. Petrol.*, 38, 201–211.