© 2001 Mineral Data Publishing, version 1.2

**Crystal Data:** Hexagonal. *Point Group:* 3m. Equant but distorted prisms, terminated by trigonal pyramids, in subparallel aggregates, to several mm.

**Physical Properties:** Fracture: Uneven to conchoidal. Tenacity: Brittle. Hardness =  $\sim$ 7 D(meas.) = 3.26(3) D(calc.) = 3.33

**Optical Properties:** Nearly opaque. *Color:* Black. *Streak:* Brown. *Luster:* Resinous, more or less splendent.

Optical Class: Uniaxial (–). Pleochroism: Strong; O = yellow-brown, dark brown, brown-black; E = dark red-brown, light brown.  $\omega = 1.800-1.820$   $\epsilon = 1.743-1.751$ 

**Cell Data:** Space Group: R3m. a = 16.186(2) c = 7.444(1) Z = 3

X-ray Powder Pattern: San Francisco mine, Bolivia. 2.63 (10), 6.63 (9), 4.05 (9), 3.05 (9), 3.61 (8), 5.13 (7), 4.71 (7)

Chemistry:

	(1)
$SiO_2$	30.7
${ m TiO}_2$	0.0
$\mathrm{B_2O_3}$	[9.0]
$\mathrm{Al_2O_3}$	1.4
$\text{Fe}_2\text{O}_3$	45.0
$V_2O_3$	$\operatorname{trace}$
FeO	1.7
MgO	6.5
$K_2O$	1.0
$\overline{\text{Na}_2}\text{O}$	2.1
$\mathrm{H_2O}$	[2.4]
Total	[99.8]

 $\begin{array}{l} (1) \ \ San \ Francisco \ mine, \ Bolivia; \ by \ electron \ microprobe, \ Fe^{2+}:Fe^{3+} \ and \ B_2O_3 \ calculated \\ from \ stoichiometry, \ H_2O \ from \ charge \ balance; \ corresponds \ to \ (Na_{0.80}K_{0.26})_{\Sigma=1.06} \\ (Fe_{2.28}^{3+}Mg_{0.53}Fe_{0.27}^{2+})_{\Sigma=3.08}(Fe_{4.29}^{3+}Mg_{1.36}Al_{0.32})_{\Sigma=5.97}(BO_3)_3Si_{5.96}O_{18}[(OH)_{3.12}O_{0.99}]_{\Sigma=4.00}. \end{array}$ 

Mineral Group: Tourmaline group.

**Occurrence:** Very rare, in fractures and lining cavities in schist metamorphosed from sedimentary rocks.

Association: Quartz, potassic feldspar, muscovite, schorl, riebeckite, magnesite.

**Distribution:** In the San Francisco mine, near Villa Tunari, Alto Chapare, Cochabamba, Bolivia.

Name: For Dr. Pavel Povondra, Charles University, Prague, Czech Republic, in recognition of work on the chemistry of the tourmaline group.

**Type Material:** Royal Ontario Museum, Toronto, Canada, M35899; National Museum of Natural History, Washington, D.C., USA, 144478.

**References:** (1) Grice, J.D., T.S. Ercit, and F.C. Hawthorne (1993) Povondraite, a redefinition of the tourmaline ferridravite. Amer. Mineral., 78, 433–436. (2) Walenta, K. and P.J. Dunn (1979) Ferridravite, a new mineral of the tourmaline group from Bolivia. Amer. Mineral., 64, 945–948.

All rights reserved. No part of this publication may be reproduced, stored in a retrieval system or transmitted in any form or by any means, electronic, mechanical, photocopying, recording, or otherwise without the prior written permission of Mineral Data Publishing.