

Steacyite**K_{1-x}(Na, Ca)₂ThSi₈O₂₀ (x = 0.2 to 0.4)**

©2001 Mineral Data Publishing, version 1.2

Crystal Data: Tetragonal. *Point Group:* 4/m 2/m 2/m. As crystals elongated || [001], terminated by {100} and {001}, to 2 mm; radiating crystalline; massive. *Twinning:* Cruciform, by 90° rotation about [010].

Physical Properties: *Tenacity:* Brittle. Hardness = 5 D(meas.) = 2.95, on porous material. D(calc.) = 3.32 Radioactive, emitting α and β particles, but not γ radiation.

Optical Properties: Translucent to transparent in thin section. *Color:* Dark brown, green, yellow. *Streak:* White. *Luster:* Vitreous.

Optical Class: Uniaxial (-). $\omega = 1.573(1)$ $\epsilon = 1.572(1)$

Cell Data: Space Group: P4/mcc. $a = 7.58(1)$ $c = 14.77(2)$ $Z = 2$

X-ray Powder Pattern: Mont Saint-Hilaire, Canada.

3.38 (100), 3.32 (55), 5.30 (45), 2.64 (41), 2.00 (26), 1.82 (20), 2.16 (19)

Chemistry:	(1)	(2)	(1)	(2)
SiO ₂	57.92	52.12	MgO	0.17
ThO ₂	28.03	30.08	CaO	4.92
UO ₂	0.00	0.14	Na ₂ O	3.36
RE ₂ O ₃	0.47	1.81	K ₂ O	3.42
Fe ₂ O ₃	0.00		F	0.23
As ₂ O ₃		0.45	H ₂ O	0.00
MnO	1.64		P ₂ O ₅	0.02
PbO	0.07		-O = F ₂	0.10
			Total	100.00
				[97.51]

(1) Mont Saint-Hilaire, Canada; corresponds to K_{0.61}(Na_{0.73}Ca_{0.72}Mn_{0.19}Mg_{0.03}) _{$\Sigma=1.67$} (Th_{0.88}Pb_{0.01}) _{$\Sigma=0.89$} Si₈O_{19.56}. (2) Rouma Isle, Guinea; RE₂O₃ = La₂O₃ 0.37%, Ce₂O₃ 0.95%, Pr₂O₃ 0.02%, Nd₂O₃ 0.38%, Sm₂O₃ 0.09%, original total given as 97.61%; corresponds to K_{0.86}(Na_{1.17}Ca_{0.72}RE_{0.10}) _{$\Sigma=1.99$} (Th_{1.04}U_{0.05}) _{$\Sigma=1.09$} (Si_{7.92}As_{0.04}) _{$\Sigma=7.96$} O₂₀.

Occurrence: In cavities in pegmatite veins in nepheline syenite in an intrusive alkalic gabbro-syenite complex (Mont Saint-Hilaire, Canada); in a miarolitic cavity in nepheline syenite (Rouma Isle, Guinea); in quartz-albite-aegirine veinlets and in albitites in syenites (Dara-i-Pioz massif, Tajikistan).

Association: Nenadkevichite, analcime, aegirine (Mont Saint-Hilaire, Canada); arfvedsonite, astrophyllite, catapleite, eudialyte, sérandite, villiaumite (Rouma Isle, Guinea); baratovite, miserite, titanite (Dara-i-Pioz massif, Tajikistan).

Distribution: From Mont Saint-Hilaire, Quebec, Canada. On Rouma Isle, Los Islands, Guinea. In the Dara-i-Pioz massif, Alai Range, Tien Shan, Tajikistan.

Name: For Harold R. Steacy, formerly Curator of the National Mineral Collection, Geological Survey of Canada, Ottawa, Canada.

Type Material: École Polytechnique, Montreal, 12480; Canadian Geological Survey, Ottawa, 61529; Canadian Museum of Nature, Ottawa, Canada; National Museum of Natural History, Washington, D.C., USA, 149820; The Natural History Museum, London, England, 1970,168.

References: (1) Perrault, G. and J.T. Szymański (1982) Steacyite, a new name, and a re-evaluation of the nomenclature of "ekanite"-group minerals. Can. Mineral., 20, 59–63. (2) (1983) Amer. Mineral., 68, 472 (abs. ref. 1). (3) Richard, P. and G. Perrault (1972) Structure cristalline de l'ekanite [steacyite] de St-Hilaire, P.Q. Acta Cryst., 28, 1994–1999 (in French). (4) Parodi, G.C. and G. Della Ventura (1987) Steacyite from the Rouma Isle (Los Islands, Republic of Guinea). Neues Jahrb. Mineral., Monatsh., 233–239. (5) Mandarino, J.A. and V. Anderson (1989) Monteregian Treasures. Cambridge Univ. Press, 188–189.

All rights reserved. No part of this publication may be reproduced, stored in a retrieval system or transmitted in any form or by any means, electronic, mechanical, photocopying, recording, or otherwise without the prior written permission of Mineral Data Publishing.